

Table 3. Texas panicum control with preemergence herbicides, Fort Cobb 2023.

Treatment	Rate		Texas Panicum Control		
			2 WAP	Peanut Injury 4 WAP	12 WAP
			%		
BAS 85001H	0.68	fl oz/a	100	100	100
BAS 85001H	1	fl oz/a	100	100	100
BAS 85001H	2	fl oz/a	100	100	100
BAS 85001H	0.68	fl oz/a	100	100	100
+ Prowl H2O	32	fl oz/a			
BAS 85001H	1	fl oz/a	100	100	100
+ Prowl H2O	32	fl oz/a			
Valor EZ	3	fl oz/a	100	100	100
+ Prowl H2O	32	fl oz/a	100	100	98
Valor EZ	3	fl oz/a	100	100	100
+ Prowl H2O	32	fl oz/a			
Valor EZ	3	fl oz/a	100	100	100
+ Pursuit	4	fl oz/a			
Valor EZ	3	fl oz/a	100	100	100
+ Prowl H2O	32	fl oz/a			
+ Pursuit	4	fl oz/a			
Valor EZ	2	fl oz/a	100	100	100
+ Prowl H2O	32	fl oz/a			
+ Pursuit	2	fl oz/a			
+ Pursuit (At Crack)	2	fl oz/a			
LSD P= .10			NS	NS	2
CV			0	0	1
All treatments applied immediately after planting unless otherwise noted. Entire trial area was treated with Select (8 fl oz/A) & 2,4-DB (16 fl oz/A) postemergence.					